

Composite label key


MDFT-Comp01

Sample type
Comp - composited
Ind - individual


Tissue type
WB - whole body
FT - fillet
CT - carcass

Species code (see legend)
Blue leader lines indicate samples collected in mudflats.

- ▲ Largemouth bass (MS)
 - ▼ Smallmouth bass (MD)
 - Brown bullhead (AN)
 - Carp (CC)
 - White catfish (AC)
 - ◆ Channel catfish (IP)
 - American eel (AR)
 - White perch (MA)
 - Northern pike (EL)
 - ◆ White sucker (WS)
- Surface sediment
- In mudflat
 - Not in mudflat
 - Bioaccumulation location (none in mudflat)
- Mudflat extent (bracketing line)
- Mudflat upper end
 - Mudflat lower end
 - Mudflat boundary
- River mile
- Sampling reach boundary
- LPRSA
- Bridge
- Abutment
- Dock



Mudflat areas for ecological receptors are those areas where the river bottom slope is $\leq 6^\circ$ and the depth is ≥ -2 ft MLLW, based on the 2007 CPG bathymetric survey. Separate mudflats were identified where there was a gap between mudflat areas or a change in the river bed which would affect mudflat formation, including tributaries, built structures, and geomorphological characteristics.





Fish samples and mudflats in the LPRSA

Reach 3

DRAFT

Privileged and Confidential: Work Product Prepared at Request of Counsel





Composite label key

MDFT-Comp01


Sample type
Comp - composited
Ind - individual

Tissue type
WB - whole body
FT - fillet
CT - carcass

Species code (see legend)
Blue leader lines indicate samples collected in mudflats.

- ▲ Largemouth bass (MS)
- ▼ Smallmouth bass (MD)
- Brown bullhead (AN)
- Carp (CC)
- White catfish (AC)
- ◆ Channel catfish (IP)
- American eel (AR)
- White perch (MA)
- Northern pike (EL)
- ◆ White sucker (WS)

- Surface sediment
- In mudflat
 - Not in mudflat
 - Bioaccumulation location (none in mudflat)
- Mudflat extent (bracketing line)
- Mudflat upper end
 - Mudflat lower end
- Mudflat boundary
- River mile
- Sampling reach boundary
- LPRSA
- Bridge
- Abutment
- Dock



Mudflat areas for ecological receptors are those areas where the river bottom slope is $\leq 6^\circ$ and the depth is ≥ -2 ft MLLW, based on the 2007 CPG bathymetric survey. Separate mudflats were identified where there was a gap between mudflat areas or a change in the river bed which would affect mudflat formation, including tributaries, built structures, and geomorphological characteristics.



Fish samples and mudflats in the LPRSA

Reach 7

DRAFT

Privileged and Confidential: Work Product Prepared at Request of Counsel

